

International Journal of Modern Physics D
 © World Scientific Publishing Company

Neutrino Oscillations at Supernova Core Bounce Generate the Strongest Gravitational-Wave Bursts

Herman J. Mosquera Cuesta

*Centro Brasileiro de Pesquisas Físicas
 CEP 22290-180 Rio de Janeiro, RJ, Brazil
 hermanjc@cbpf.br*

Karen Fiuza

*Instituto de Física, Universidade Federal do Rio Grande do Sul
 CEP 91501-970 Porto Alegre, RS, Brazil
 kfiuza@if.ufrgs.br*

Received 18 December 2003

Revised 23 January 2004

During the core bounce of a supernova collapse resonant active-to-active ($\nu_a \rightarrow \nu_a$), as well as active-to-sterile ($\nu_a \rightarrow \nu_s$) neutrino (ν) oscillations can take place. Besides, over this phase weak magnetism increases antineutrino ($\bar{\nu}$) mean free paths, and thus its luminosity. Because the oscillation feeds mass-energy into the target ν species, the large mass-squared difference between species ($\nu_a \rightarrow \nu_s$) implies a huge amount of power to be given off as gravitational waves ($L_{\text{GWs}} \sim 10^{49} \text{ erg s}^{-1}$), due to anisotropic but coherent ν flow over the oscillation length. This anisotropy in the ν -flux is driven by both the *universal spin-rotation* and the spin-magnetic coupling. The new spacetime strain estimated this way is still several orders of magnitude larger than those from ν diffusion (convection and cooling) or quadrupole moments of the neutron star matter. This new feature turns these bursts the more promising supernova gravitational-wave signal that may be detected by observatories as LIGO, VIRGO, etc., for distances far out to the VIRGO cluster of galaxies.

Keywords: elementary particles: neutrino; gravitational waves; supernova

1. Generating GWs During ν Oscillations

To start with, let us recall how the production of GWs during ν oscillations proceeds by considering the case of oscillations between active and sterile neutrinos in the supernova core. The essential point here is that oscillations into sterile neutrinos change dramatically the energy and momentum (linear and angular) configuration of the proto-neutron star (PNS). In particular, flavor conversions into sterile neutrinos drive a large mass and energy loss from the PNS because once they are produced they freely escape from the star. This means that oscillations into sterile, in dense matter, take place over longer oscillation lengths, compared to $\nu_a \rightarrow \nu_a$, and the sterile encounter infinite mean free paths thereafter. Physically, the potential, $V_s(x)$,

for sterile neutrinos in dense matter is zero, and their probability of reversion, still inside the star, into active species is quite small. This outflow translates into a noticeable modification of the PNS mass and energy quadrupole

Since most sterile neutrinos escape along the directions defined by the dipole field and angular momentum vectors (see Fig.2), then the ν outflow is at least quadrupolar in nature. This produces a super strong gravitational-wave burst once the flavor conversions take place, the energy of which stems from the energy and momentum of the total number of neutrinos participating in the oscillation process^a.

The remaining configuration of the star must also reflect this loss, i.e., the active ν s were coupled to the neutron matter. Hence its own matter and energy distribution becomes also quadrupolar. Because this quadrupole configuration (the matter and energy still trapped inside the just-born neutron star) keeps changing over the timescale for which most of the oscillations take place, then GWs must also be emitted from the star over that transient. At the end, the probability of conversion and the ν -flux anisotropy parameter (α , see below) determine both how much energy partakes in the process and the degree of asymmetry during the emission.

The case for oscillations among active species is a bit different, the key feature being that mass and energy is coherently relocated (see Ref.¹) from one region to another inside the PNS, especially because of the *weak magnetism* of antineutrinos that allows them to have larger mean free paths². In addition, oscillations of electron neutrinos into muon or tauon neutrinos leave these last species outside their own neutrino spheres, and hence they are in principle free to stream away. This must also generate GWs during that sort of flavor conversions, although their specific strength (strain) must be a bit lower compared to conversions into sterile neutrinos wherein almost all the ν species may participate, and the large Δm^2 in the process. These neutral-current interacting ν species must be the very first constituents of the ν burst from any supernova since most ν_e s are essentially trapped.

Since the sterile neutrinos escape the core over a timescale of a few milliseconds (ms), the number of neutrinos escaping and their angular distribution is sensitive to the instantaneous distribution of neutrino production sites. Since thermalization cannot occur over such short times ($\Delta T^{\text{thermal}} \sim 0.5$ s), and since the neutrino production rate is sensitive to the local temperature at the production site, the inhomogeneities during the collapse phase get reflected in the inhomogeneities in the escaping neutrino fluxes and their distributions. Because of both the ν spin- $\vec{B}$ and ν spin- $\vec{J}$ coupling the asymmetries in these distributions can give rise to quadrupole moments, which must generate gravitational waves^{3,4}, and dipole moments which can explain the origin of pulsar kicks^{5,6}.

Fixed by the *probability of oscillation*, $P_{\nu_a \rightarrow \nu_a}$, the *fraction* of neutrinos that can escape in the first few milliseconds is, however, *small*. Firstly, the neutrinos have to be produced roughly within one mean free path from their resonance surface.

^aThe attentive reader must regard that neutrinos carry away almost all of the binding energy of the just-born neutron star, i.e., $\Delta E_\nu \sim 3 \times 10^{53}$ erg.

Secondly, since in the case of $\nu_a \rightarrow \nu_s$ oscillations the ν_s is the heaviest neutrino species, the sign of the effective potential $A(x)$ and the resonance condition indicates that only ν_e s and the antineutrinos $\bar{\nu}_\mu$ and $\bar{\nu}_\tau$ can undergo resonant conversions.

2. Anisotropic Neutrino Outflow: origin and computation

To provide a physical foundation for the procedure introduced here to determine the neutrino *asymmetry parameter*, α , which measures how large is the deviation of the ν -flux from a spherical one, we recall next two fundamental effects that run into action once a PNS is forming after the SN collapse. We stress in passing that other physical process such as convection, thermalization, etc., are not relevant over the timescale under consideration: $\Delta T_{\text{osc}} \sim 3 - 10$ ms after the SN core bounce, and therefore do not modify in a sensitive manner the picture described below. Indeed, if ν thermalization, for instance, already took place, then the oscillations are severely precluded since oscillations benefit of the existence of energy, matter density and entropy gradients inside the PNS^{7,8}, which are “erased” once thermalization onsets.

As discussed by Mosquera Cuesta and Fiuza¹ (see also the detailed discussion by Loveridge⁶ about the ν distribution function inside the PNS, an asymmetric, off-centered rotating ν beam) both the ν -spin coupling to rotation and the ν -spin coupling to the magnetic ($\vec{B}$) field drives an effective momentum, and thus energy flux, asymmetry along the angular momentum ($\vec{J}$) and $\vec{B}$ -field directions. The combined action on the escaping ν s of a rotating background spacetime plus magnetic field makes their ν -sphere a decidedly distorted surface, with at least a quadrupolar distribution. This is the source of the strong GWs burst in this mechanism when the oscillations ensue.

Based on the concomitant action of both the effects: ν -spin coupling to both the magnetic field and rotation, described previously, one can determine the ν flow anisotropy in a novel, self-consistent fashion by defining α as the ratio of the total volume filled by the distorted ν -spheres to that of the proto-neutron star (PNS). The ν_e -sphere radius of a non-magnetic non-rotating star is obtained from the condition:

$$\tau_{\nu_e}(R_{\nu_e}) = \int_{R_{\nu_e}}^{\infty} \mathcal{K}_{\nu_e} \rho(r) dr = \frac{2}{3},$$

where τ_{ν_e} is the optical depth, and $\mathcal{K}_{\nu_e}$ the scattering opacity for electron neutrinos, and ρ the matter density.

One can take hereafter⁹ $R_{\text{PNS}}^{\nu_e} \equiv R_\nu \sim 35$ km, which is of the order of magnitude of the oscillation length λ_ν of a typical supernova ν ; which is constituent as well of the atmospheric ν s, for which $\Delta m^2 \sim 10^{-3} \text{eV}^2$ has been estimated by Superkamiokande ν detector¹⁰

$$\lambda_\nu \sim 31 \text{ km} \left[\frac{E_{\nu_e}}{10 \text{ MeV}} \right] \left(\frac{10^{-3} \text{ eV}^2}{\Delta m^2} \right).$$

Therefore, resonant conversions between active species may take place at the posi-

110 *H. Mosquera Cuesta, K. Fiuza*

tion r from the center defined by

$$r = R_{\nu_e} + \delta_0 \cos \phi ,$$

with ϕ the angle between the ν -spin and $\vec{B}$, i.e.,

$$\cos \phi = \frac{(\vec{k} \cdot \vec{B})}{k} ; \delta_0 = \frac{e\mu_e B}{2\pi^2(dN_e/dr)} \sim 1 - 10 \text{ km} ,$$

for $B \sim 10^{14-15}$ G, respectively. Here e , $N_e = Y_e N_n$ and μ_e represent the electron charge, density and chemical potential, respectively.

This defines in Fig.2 an ellipsoidal figure of equilibrium with semi axes $a = R_{\nu} + \delta_0$; and $b = R_{\nu}$, and volume (after rotating around $\vec{B}$):

$$V_{ellips.} = \frac{4}{3}\pi R_{\nu}^2(R_{\nu} + \delta_0) .$$

Meanwhile, the ν -spin coupling to rotation generates an asymmetric lemniscate-like plane curve (see Fig.2) $r = R_{\nu} (1 \pm L a \cos \theta)^{1/2}$, which upon a 2π rotation around the star angular momentum axis generates a volume:

$$V_{lemnisc.} = \frac{1}{4}(2R_{\nu})^2 \times 2\pi \bar{y} + \frac{1}{4}(R_{\nu})^2 \times 2\pi \bar{y} ,$$

where the quantity $\bar{y}$ (and $\bar{x}$) is defined as the location of the centroid of one of the lobes of that plane figure with respect to its coordinate center (x,y). After a long, but straightforward, calculation one obtains $\bar{y} = \frac{\sqrt{\pi}}{\Gamma^2(1/4)} R_{\nu}$, $\bar{x} = \frac{4\sqrt{\pi}}{\Gamma^2(1/4)} R_{\nu}$. Thence, for a PNS⁹ with parameters cited in this article one obtains¹¹

$$\alpha_{\min} = \frac{V_{ellips.} + V_{lemnisc.}}{V_{\text{PNS}}} \sim 0.11 - 0.01 . \quad (1)$$

As such, this is essentially a new result of this paper. The attentive reader must notice in passing that the definition in Eq.(1) does take into account all of the physics of the neutrino oscillations: luminosity, density gradients and angular propagation, since Fig.2 does gather the relevant information regarding the spatial configuration of the ν luminosity in as much as is done in the standard definition of α ^{9,12}

$$\alpha(t) \equiv \frac{1}{L_{\nu}(t)} \int_{4\pi} d\Omega' \Psi(\theta', \phi') \frac{L_{\nu}(\Omega', t)}{d\Omega'} . \quad (2)$$

Indeed, one can get the "feel" of the relationship between these two definitions by noticing that the quantity $L_{\nu}(\Omega', t)$ in the integrand of Eq.(2) can be expressed as $L_{\nu}(\Omega', t) \equiv L_{\nu}(t)F(\Omega')$, where the function $F(\Omega')$ contains now all the information regarding the angular distribution of the neutrino emission. Hence $L_{\nu}(t)$ can be factorized out of the integral and dropped from Eq.(2). This converts Eq.(2) in a relationship among (*solid*) angular quantities, which clearly can be reduced to a volumetric one, similar to the one introduced in Eq.(1), upon a transformation using the definition of *solid angle* in the form of *Lambert's law*: $d\Omega = \frac{dA \cos \theta}{R^2}$, and applied to the sphere representing the PNS. Here θ , measured from a coordinate

system centered at the PNS¹², plays the role of the angle between the direction towards the observer and the direction Ω' of the radiation emission in Eq.(24) of Ref.¹². Therefore, the novel result here presented is physically consistent with the standard one for the asymmetry parameter α .




Fig. 1. A schematic representation of the ν -flux distribution (hatched regions) in a nascent pulsar. The at least quadrupolar distribution is evident and stems from the neutrino spin-magnetic and spin-rotation couplings.

3. Enlarged ν and GWs Luminosity From Oscillations in Dense Matter

Numerical simulations¹² showed that, in general, the fraction of the total binding energy emitted as GWs by pure ν convection^{3,11,12} is: $E_{\text{GW}}^{\nu} \sim [10^{-10}-10^{-13}] M_{\odot} c^2$, for a ν total luminosity: $L_{\nu} \sim 10^{53} \text{ erg s}^{-1}$. In the production of GWs via ν oscillations³ ($\nu_a \leftrightarrow \nu_a$ or $\nu_a \leftrightarrow \nu_s$) there exists two main reasons for expecting a major enhancement in the GWs luminosity during the transition: a) the conversion itself, which makes the overall luminosity (L_{ν_x}) of a given ν_x species to grow by a large factor: $\Delta L_{\nu_x} \lesssim 10\% L_{\nu}^{\text{total}}$. The enhancement stems from the mass-energy being given to, or drained from, the new ν species into which oscillations take place. This augment gets reflected in the species mass-squared difference Δm^2 , and their relative abundances: one species is number-depleted while the other gets its number enhanced.

But, even if the energy increase, or give up, is small, b) the abrupt (resonant) conversion over the transition time $\Delta T_{\text{osc}} \equiv \frac{\lambda_{\text{osc}}}{V_{\nu-Diff.}} \sim [10^{-4} - 10^{-3}]$ s, also magnifies transiently L_{ν_e} . Here λ_{osc} defines the oscillation length, and $\bar{V}_{\nu-Diff.} \sim 10^9 \text{ cm s}^{-1}$ the convective ν diffusion velocity¹².

If flavor transitions can indeed take place during supernovae (SNe) core collapse and bounce, then they must leave some imprints in the SNe neutrino spectrum. Main observational consequences of neutrino conversions inside SNe include the partial or total disappearance of the neutronization peak; the moment at which most ν_e s are produced, the interchange of the original spectrum and the appearance of a hard ν_e spectrum, together with distortions of the ν_e energy spectrum and alterations of the $\bar{\nu}_e$ spectrum¹³.

3.1. *Resonance, Adiabaticity and the Role of Weak Magnetism*

ν oscillations in vacuum produce no GWs^{3,4}. In the case of active-to-active ν oscillations (essentially the same physical argument holds also for active-to-sterile ν oscillations), the main reason for this negative result is that this class of conversions do not increase in a significant figure the total number of particles escaping from the proto-neutron star. In dense matter, the process generates no GWs since the oscillations develop with the neutrinos having very short mean free paths, so that they motion can be envisioned as a standard diffusion process.

However, because of the active *antineutrino* species *weak magnetism* their effective luminosity can be enlarged as much as 15% compared to the typical one they achieve when this effect is not taken into consideration during their propagation in dense matter². This result can be interpreted by stating that the number of oscillating (and potentially escaping) antineutrinos may be augmented by a large factor because now they do encounter longer mean free paths. Below we take advantage of this peculiar behavior of the ν outflow in supernovae in computing the overall probability of transition between active species.

3.2. *Active-To-Active ν Oscillations*

The neutrino oscillation pattern in vacuum can get noticeably modified by the passage of neutrinos through matter because of the effect of *coherent forward scattering*^{14,15}. Therefore, interaction with matter may help in allowing more ν s to escape if resonant conversions into active³ and/or sterile ν s³ occur inside the ν -sphere of the active ν s. The transition probability⁷ for conversions among active species thus reads

$$P_{\nu_a \rightarrow \nu_a}(|\vec{x}_1 - \vec{x}_0|) = 1 - \frac{1}{2} [1 + \cos 2\theta_m(x_0) \cos 2\theta_m(x_1)], \quad (3)$$

where x_0 and x_1 correspond to the production and detection sites, respectively, and θ_m the mixing angle in matter.

In order to produce an effect, neutrinos must be able to escape the core without thermalizing with the stellar material. For active neutrino species of energies ≈ 10

MeV, this is not possible as long as the matter density is $\gtrsim 10^{10} \text{ g cm}^{-3}$. Since the production rate of neutrinos is a steeply increasing function of matter density (production rate $\propto \rho^n$, where ρ is the matter density and $n > 1$), the overwhelming majority of the neutrinos of all species produced are trapped. This way, there seems to be no contribution to the GWs amplitude for neutrino conversions taking place within the active neutrino flavors. Nonetheless, if there were indeed *weak magnetism effects*², one can rethink of conversions within active species. In the new result on *weak magnetism for antineutrinos in core collapse supernovae*, the antineutrinos ($\bar{\nu}_x$ s) luminosity could be noticeably increased because of their longer mean free paths², and this means that the total energy flux can be augmented in $\sim 15\%$ for ν s of temperature $\sim 10 \text{ MeV}$. One can verify that longer mean free paths allows for a larger oscillation probability, and hence the contribution to the generation of GWs during flavor conversions within active species becomes non-negligible compared to the earlier case³ where not weak magnetism effects were taken into account.

For active-to-active ν flavor conversions, for instance: $\nu_e \rightarrow \nu_\mu, \nu_\tau$; as implied by SNO results, the resonance must take place at a distance x_{res} from the PNS center and amid the active ν -spheres, whenever the resonance condition is satisfied

$$4\Delta m^2 \cos 2\theta_m = 2\sqrt{2} G_F N_e(x_{\text{res}}) k_{\nu_e};$$

notice that we neglected the magnetic field contribution. Here $k_{\nu_e} = E_{\nu_e}/c$ is the ν_e momentum, and $N_e(x_{\text{res}}) = N_{e-} - N_{e+} \sim 10^{39-40} \text{ cm}^{-3}$ is the electron number density. Thus the right-hand part of this equation reduces to

$$2\sqrt{2}G_F E_{\nu_e} N_e(x_{\text{res}}) = 6.9 \times 10^4 \text{ eV}^2 \left[\frac{\rho}{10^{11} \text{ g cm}^{-3}} \right] \left(\frac{E_{\nu_e}}{10 \text{ MeV}} \right). \quad (4)$$

For densities of order $\rho_{x=x_{\text{res}}} \sim 10^{10} \text{ g cm}^{-3}$, i.e., recalling that ν_e s are produced at the PNS outermost regions³ where the electron to baryon ratio is $Y_e \sim 0.1$, the resonance condition is satisfied for $\Delta m^2 \sim 10^4 \text{ eV}^2$, which implies a neutrino mass of about $m_\nu \sim 10^2 \text{ eV}$. Neutrino flavor conversions in the resonance region can be strong if the adiabaticity condition is fulfilled³, i.e., whenever⁷ $\frac{\Delta m^2 \sin^2 2\theta_m}{2E_\nu \cos 2\theta_m} \left(\frac{1}{\rho} \frac{d\rho}{dx} \right)^{-1}_{x=x_{\text{res}}} \gg 1$, where x_{res} is the position of the resonance layer. Recalling that the typical scale of density variations in the PNS core is $h_{N_e} \sim (dN_e/dr)^{-1} \sim 6 \text{ km}$, this adiabatic behavior could be achieved as far as the density and magnetic field remain constant over the oscillation length

$$\lambda_{\text{osc}} \equiv \left(\frac{1}{\rho} \frac{d\rho}{dx} \right)^{-1}_{x=x_{\text{res}}} \sim \left(\frac{1}{2\pi} \frac{\Delta m^2}{2k_\nu} \sin(2\theta_m) \right)^{-1} \sim \frac{1 \text{ cm}}{\sin(2\theta_m)}, \quad (5)$$

of order h_{N_e} , which can be satisfied for $\Delta m^2 \sim 10^4 \text{ eV}^2$ as long as¹⁹ $\sin^2 2\theta_m > 10^{-8}$. Although these ν oscillations could be adiabatic for a wide range of mixing angles, and thus a large number of ν s could actually oscillate, a ν_x mass such as this is incompatible with both viable solutions to the Solar Neutrino Problem (SNP) and the most recent cosmological constraints on the total mass of all stable neutrino species that could have left their imprint in the Cosmic Microwave Background

Radiation (CMBR), as inferred from the observations performed by the satellite WMAP: $m_\nu \sim 1$ eV. Therefore, we dismiss this possibility since there appears to be no evidence for neutrinos masses in this parameter range inside the PNS core.

On the other hand, if one takes into account the KamLAND results¹⁶, which definitively demonstrated that *i*) a large mixing angle (LMA) solution of the solar ν problem is favored: $\sin^2 2\theta \sim 0.8$, *ii*) for a mass-squared difference: $\Delta m^2 \sim 5.5 \times 10^{-5}$ eV² (we use the approximate value $\Delta m^2 \sim 10^{-4}$ eV² for the estimates below), one can see that resonant conversions with $\Delta m^2 = m_{\nu_2}^2 - m_{\nu_1}^2 \sim 10^{-4}$ eV² would take place in supernova regions where the density is about $\rho \sim 10 - 30$ g cm⁻³, which corresponds to the outermost layers of the exploding star. Although a large number of ν species can in effect participate of the transitions there, i.e, the neutrino luminosity can be still a very large quantity, these regions are of no interest for the gravitational-wave emission from neutrino oscillations since the overall energy density at that distance from the star center is very small. This does not mean that no GWs are emitted from transitions there, it is to mean that their strain is very small so as to be detectable. Observations of ν oscillations¹³ in this parameter range would provide useful information regarding the SNP, the hierarchy of neutrino masses and the mixing $|U_{e3}|^2$. Note in passing that oscillations in this range would imply a mass for the ν_2 species $m_{\nu_2} \sim 10^{-2}$ eV ($\frac{\rho}{10 \text{ g cm}^{-3}}$), in the case when $m_{\nu_2} > m_{\nu_1}$. This m_{ν_2} is compatible with current limits from WMAP²⁰.

Finally, let us consider oscillations in the parameter range estimated from CMBR by WMAP observations. In this case, resonant ν transitions would take place in regions where the density is as high as $\rho \sim 10^{7-8}$ g cm⁻³, that is, at the supernova mantle or PNS upper layers. At these densities the oscillation length can be still $\lambda_{\text{osc}} \sim 1 - 5$ km, and thus the conversions can be considered as adiabatic. Thus the resonance condition can be satisfied for $\Delta m^2 \sim 1$ eV² as long as $\sin^2 2\theta_m \lesssim 10^{-3}$.

Hence, by taking with this constraint, and recalling that *i*) at least 6 ν -species can participate in the flavor transitions, *ii*) most ν s are emitted having $\vec{k}_\nu$ parallel to $\vec{B}$, which implies a further reduction factor of 2, and *iii*) most ν s are emitted having $\vec{k}_\nu$ parallel to $\vec{J}$ implying also an additional reduction factor of 2, one can show from Eq.(3) that the *fraction* of ν_a species that can eventually exchange flavor during the first few milliseconds after core bounce turns out to be $P_{\nu_a \rightarrow \nu_a}(|\vec{x} - \vec{x}_{res}|) \lesssim 0.1\%$. Thus, the total energy involved in the oscillation process we are considering could be estimated as: $E_\nu^{\text{tot}} \simeq P_{\nu_a \rightarrow \nu_a}(|\vec{x} - \vec{x}_{res}|) \times N_\nu k_\nu c (1 + 0.15)$, with N_ν the total number of neutrinos undergoing flavor conversions during the timescale ΔT_{osc} . Note in passing that KamLAND $\bar{\nu}_e$ experiment suggests¹⁶ $P_{\bar{\nu}_e \rightarrow \bar{\nu}_{\mu,\tau}} \sim 40\%$, while LSND $P_{\bar{\nu}_\mu \rightarrow \bar{\nu}_e} \sim 0.26\%$. This result from LSND has not been so far ruled out by any terrestrial experiment, and there is a large expectation that it could be verified by MiniBoone at Fermilab.

In the case of active-to-sterile ν conversions, it was proved in Refs.^{3,4,1} (see details therein) that a substantial fraction: $P_{\nu_a \rightarrow \nu_s}$, of ν s may get converted to

sterile ν_s , and escape the core of the star, if the sterile ν mass (m_{ν_4}) is such that

$$1 \text{ eV}^2 \lesssim \Delta m_{\nu_a \rightarrow \nu_4}^2 \lesssim 10^4 \text{ eV}^2. \quad (6)$$

For both classes of ν conversions the number of ν_s escaping and their angular distribution is sensitive to the instantaneous distribution of production sites. These inhomogeneities can give rise to quadrupole moments that generates GWs^{3,12}, and dipole moments that could drive the runaway pulsar kicks^{5,6}. Noting that at least 6 ν -species can participate in both types of oscillations and that the interaction with both the magnetic field and angular momentum of the PNS brings with an overall reduction factor of 4 in the oscillation probability, one can show that the *fraction* of ν_s that can actually undergo flavor transitions in the first few milliseconds is³

$$P_{\nu_a \rightarrow \nu_s}^{\nu_a \rightarrow \nu_a}(|\vec{x} - \vec{x}_0|) \lesssim 1\%, \quad (7)$$

of the total ν_s number: $M_\odot \times m_p^{-1} \sim 10^{57} \nu$, which corresponds to a total energy exchanged during the transition

$$\Delta E_{\nu_a \rightarrow \nu_s}^{\nu_a \rightarrow \nu_a} \sim 3 \times 10^{51} \text{ erg}. \quad (8)$$

Equations (8,3) determine the total luminosity of the neutrinos participating in the resonant flavor conversions.

4. GWs Energetics From ν Luminosity and Detectability

If ν oscillations do take place in the SN core, then the most likely detectable GWs signal should be produced over the time interval for which the conditions for flavor conversions to occur are kept, i.e., $\sim (10^{-1} - 10)$ ms. This timescale implies GWs frequencies in the band: $f_{\text{GWs}} \sim [10 - 0.1]$ kHz, centered at 1 kHz, because of the maximum ν production around 1 – 3 ms after core bounce³. This frequency range includes the optimal ban width for detection by ground-based observatories. For a 1 ms conversion time span the ν luminosity reads

$$L_\nu \equiv \frac{\Delta E_{\nu_a \rightarrow \nu_s}}{\Delta T_{\text{osc}}} \sim \frac{3 \times 10^{51} \text{ erg}}{1 \times 10^{-3} \text{ s}} = 3 \times 10^{54} \frac{\text{erg}}{\text{s}}. \quad (9)$$

To estimate the GWs burst amplitude produced by the non-spherical outgoing front of oscillation-produced ν_s one can write (see details in Refs.^{1,3,11,12})

$$h_\nu = \frac{2G}{c^4 R} (\Delta t \times L_\nu \times \alpha) \simeq |A| \left[\frac{55 \text{ kpc}}{R} \right] \left(\frac{\Delta T}{10^{-3} \text{ s}} \right) \left[\frac{L_\nu}{3 \times 10^{54} \frac{\text{erg}}{\text{s}}} \right] \left(\frac{\alpha}{0.1} \right),$$

where $|A| = 4 \times 10^{-23} \text{ Hz}^{-1/2}$ is the amplitude.^b A GWs signal this strong will likely be detected by the first generation of GWs interferometers as LIGO, VIRGO,

^bOne must realize at this point that the high value of the anisotropy parameter here used is consistently supported by the discussion by Mosquera Cuesta and Fiuza¹ regarding the neutrino coupling to rotation and magnetic field quoted above.

etc. Its imprint in the GWs waveform may resemble a spike of high amplitude and time width of $\sim$ ms followed by a Christodoulou's memory⁴. From Eq.(10) the GWs luminosity turns out to be

$$L_{\text{GWs}} \sim 10^{48-50} \frac{\text{erg}}{\text{s}} \left[\frac{L_\nu}{3 \times 10^{54} \frac{\text{erg}}{\text{s}}} \right]^2 \left(\frac{\alpha}{0.1 - 0.01} \right)^2, \quad (10)$$

while the GWs energy radiated in the process yields $E_{\text{GWs}} \equiv L_{\text{GWs}} \times \Delta T_{\text{osc}} \sim 10^{47-45} \text{ erg}$. This is about $10^{5-3} \times L_\nu^{\text{Diff}}$ the luminosity from ν diffusion inside the PNS⁴.

5. Conclusion

One can see that if ν flavor conversions indeed take place during SN core bounce inasmuch as they take place in our Sun and Earth²³, then GWs should be released during the transition. The GWs signal from the process is expected to irradiate much more energy than current mechanisms figured out to drive the NS dynamics at birth do. A luminosity this large (Eq.(9)) would turn these bursts the strongest GWs signal to be detected from any SN that may come to occur on distances up to the VIRGO cluster, $R \sim [10 - 20]$ Mpc. It is stressed that this signal will still be the stronger one from a given SN, even in the worst case in which the probability of ν conversion is three orders of magnitude smaller than the estimated in the present paper. In proviso, we argue that a GWs signal that strong could have been detected during SN1987a from the Tarantula Nebula in the Milky Way's satellite galaxy Large Magellanic Cloud, despite of the low sensitivity of the detectors at the epoch. In such a case, the GWs burst must have been correlated in time with the earliest arriving neutrino burst constituted of some active species given off during the very early oscillation transient where some ν_e s went into ν_μ s, ν_τ s or ν_s s. Thenceforth, it could be of worth to reanalyze the data collected far from that event taking careful follow up of their arrival times, if appropriate timing was available at that moment.

6. Acknowledgements

The authors are indebted to Prof. César Vasconcellos for inviting us to participate in this important meeting, which make it possible to start with this our collaboration. HJMC acknowledges the support from FAPERJ, Brazil, through the grant E-26/151.684/2002. KF thanks CAPES (Brazil) for a graduate fellowship.

References

1. H. J. Mosquera Cuesta and K. Fiuza, to be published in *Eur. Phys. J. C* (2004).
2. C. J. Horowitz, *Phys. Rev.* **D65** (2002) 043001.
3. H. J. Mosquera Cuesta, *ApJ* **544** (2000) L61.
4. H. J. Mosquera Cuesta, *Phys. Rev. D* **65** (2002) 061503(R).
5. G. M. Fuller, A. Kusenko, I. Mocioiu & S. Pascoli, *Phys. Rev.* **D68** (2003) 103002.
6. L. C. Loveridge, *Phys. Rev.* **D69** (2004) 024008.

7. W. Grimus, *Neutrino Physics - Theory* hep-ph/0307149 (2003).
8. E. K. Akhmedov et al., *Nuc. Phys.* **B542** (1999) 3.
9. A. Burrows, J. Hayes & B. A. Fryxell, *ApJ* **450** (1995) 830.
10. Y. Fukuda et al., *Phys. Rev. Lett.* **81** (1998) 1562.
11. A. Burrows & J. Hayes, *Phys. Rev. Lett.* **76** (1996) 352.
12. E. Müller & H.-Th. Janka, *A&A* **317** (1997) 140.
13. A. S. Dighe & A. Yu. Smirnov, *Phys. Rev.* **D62** (2000) 033007.
14. L. Wolfenstein, *Phys. Rev.* **D20** (1979) 2634.
15. S. P. Mikheyev & A. Yu. Smirnov, *Sov. J. Nucl. Phys.* **42** (1985) 913.
16. K. Eguchi et al. [KamLAND Collaboration], *Phys. Rev. Lett.* **90** (2003) 021802.
17. H.-Th. Janka, K. Kifonidis & M. Rampp, astro-ph/0103015 (2001).
18. A. Vilenkin, *Phys. Rev. Lett.* **41** (1978) 1575.
19. A. Kusenko, private communication. Also A. Kusenko, astro-ph/9903167.
20. A. Pierce & H. Murayama, *New neutrino mass constraints based on the Wilkinson Microwave Anisotropy Probe (WMAP)* hep-ph/0302131.
21. A. Kusenko & G. Segrè, *Phys. Lett.* **B396** (1997) 197.
22. H. Nunokawa et al., *Phys. Rev.* **D56** (1997) 1704.
23. A. Yu. Smirnov, *Nuov. Cim.* **B117** (2002) 1237.